

available at [www.sciencedirect.com](http://www.sciencedirect.com)

SciVerse ScienceDirect

journal homepage: [www.elsevier.com/locate/jjcc](http://www.elsevier.com/locate/jjcc)

## Volumes Contents

### VOLUME 57, ISSUE 1

January 2011

#### Editorial

- It's Time for the Real International Journal. *A. Hirayama* . . . . . 1

#### Reviews

- The role of vascular failure in coronary artery spasm. *H. Kawano, K. Node* . . . . . 2
- Chronic kidney disease and heart failure—Bidirectional close link and common therapeutic goal.  
*N. Shiba, H. Shimokawa* . . . . . 8
- Recent advances in coronary angiography. *Y. Uchida* . . . . . 18
- Mild to moderate aortic stenosis and coronary bypass surgery. *X. Du, J.L. Soon* . . . . . 31

#### Original Articles

- Cavity volume of ruptured plaque is an independent predictor for angiographic no-reflow phenomenon during primary angioplasty in patients with ST-segment elevation myocardial infarction. *K. Ohshima, S. Ikeda, H. Kadota, K. Yamane, N. Izumi, H. Kawazoe, K. Ohshima, M. Hamada* . . . . . 36
- Retrospective comparison of clinical and angiographic outcomes after primary stenting using sirolimus-eluting and bare-metal stents in nonrandomized consecutive 568 patients with first ST-segment elevated myocardial infarctions. *T. Ishikawa, M. Mutoh, Y. Nakano, A. Endo, T. Kubota, T. Suzuki, K. Nakata, A. Murakami, T. Miyamoto, H. Sakamoto, H. Okada, K. Imai, M. Yoshimura* . . . . . 44
- What is the optimal duration of triple anti-platelet therapy in patients with acute myocardial infarction undergoing drug-eluting stent implantation? *K.-H. Park, M.H. Jeong, M.G. Lee, J.S. Ko, D.S. Sim, N.S. Yoon, H.J. Yoon, Y.J. Hong, K.H. Kim, H.W. Park, J.H. Kim, Y. Ahn, J.G. Cho, J.C. Park, J.C. Kang* . . . . . 53
- Relationship between renal function stage and clinical outcomes after paclitaxel-eluting stent implantation. *Y. Yazaki, R. Iijima, M. Nakamura, K. Sugi* . . . . . 61
- Prevalence of anomalous origin of coronary artery detected by multi-detector computed tomography at one center. *S. Fujimoto, T. Kondo, T. Orihara, J. Sugiyama, M. Kondo, T. Kodama, H. Fukazawa, H. Nagaoka, A. Oida, J. Yamazaki, S. Takase* . . . . . 69
- Long-term event monitoring study of fluvastatin in Japanese patients with hypercholesterolemia: Efficacy and incidence of cardiac and other events in elderly patients ( $\geq 65$  years old). *H. Itakura, N. Nakaya, T. Kusunoki, N. Shimizu, S. Hirai, S. Mochizuki, T. Ishikawa* . . . . . 77

Impact of reduced left atrial functions on diagnosis of paroxysmal atrial fibrillation: Results from analysis of time-left atrial volume curve determined by two-dimensional speckle tracking. <i>M. Mori, H. Kanzaki, M. Amaki, T. Ohara, T. Hasegawa, H. Takahama, K. Hashimura, T. Konno, K. Hayashi, M. Yamagishi, M. Kitakaze</i> . . . . .	89
Investigation of optimal anticoagulation strategy for stroke prevention in Japanese patients with atrial fibrillation—The J-RHYTHM Registry study design. <i>H. Atarashi, H. Inoue, K. Okumura, T. Yamashita, H. Origasa, for the J-RHYTHM Registry Investigators</i> . . . . .	95
Improvement of autonomic nervous activity by Waon therapy in patients with chronic heart failure. <i>S. Kuwahata, M. Miyata, S. Fujita, T. Kubozono, T. Shinsato, Y. Ikeda, S. Hamasaki, T. Kuwaki, C. Tei</i> . . . . .	100
Predicting the success of noninvasive positive pressure ventilation in emergency room for patients with acute heart failure. <i>A. Shirakabe, N. Hata, S. Yokoyama, T. Shinada, N. Kobayashi, K. Tomita, M. Kitamura, A. Nozaki, H. Tokuyama, K. Asai, K. Mizuno</i> . . . . .	107
Promoter-specific lentivectors for long-term, cardiac-directed therapy of Fabry disease. <i>C.-J. Lee, X. Fan, X. Guo, J.A. Medin</i> . . . . .	115
Common knowledge regarding prevention of infective endocarditis among general dentists in Japan. <i>K. Nakano, T. Ooshima</i> . . . . .	123

## **VOLUME 57, ISSUE 2 . . . . . March 2011**

### **Reviews**

Current biochemistry, molecular biology, and clinical relevance of natriuretic peptides. <i>T. Nishikimi, K. Kuwahara, K. Nakao</i> . . . . .	131
Granzyme B as a novel factor involved in cardiovascular diseases. <i>Y. Saito, H. Kondo, Y. Hojo</i> . . . . .	141
Heart failure due to abnormal filling function of the heart. <i>D.-W. Sohn</i> . . . . .	148

### **Original Articles**

Associations of positive T wave in lead aVR with hemodynamic, coronary, and left ventricular angiographic findings in anterior wall old myocardial infarction. <i>K. Shinozaki, A. Tamura, J. Kadota</i> . . . . .	160
Atypical presentation of acute coronary syndrome: A significant independent predictor of in-hospital mortality. <i>A. El-Menyar, M. Zubaid, K. Sulaiman, W. AlMahmeed, R. Singh, A.A. Alsheikh-Ali, J. Al Suwaidi, for the Gulf Registry of Acute Coronary Events (Gulf RACE) Investigators</i> . . . . .	165
Impact of early statin initiation on secondary prevention in Japanese patients with coronary artery disease. <i>M. Nakamura, T. Yamashita, J. Yajima, Y. Oikawa, K. Ogasawara, K. Sagara, A. Koike, H. Kirigaya, K. Nagashima, T. Otsuka, T. Uejima, R. Funada, S. Matsuno, S. Suzuki, H. Sawada, T. Aizawa, the Shinken Database Study Group</i> . . . . .	172
Effect of preoperative hemoglobin A1c levels on long-term outcomes for diabetic patients after off-pump coronary artery bypass grafting. <i>R. Tsuruta, K. Miyauchi, T. Yamamoto, S. Dohi, K. Tambara, T. Dohi, H. Inaba, K. Kuwaki, H. Daida, A. Amano</i> . . . . .	181
Angiographic late lumen loss at the site of overlap of multiple Cypher™ sirolimus-eluting stents: ALSOCE study. <i>Y. Takamiya, S.-i. Miura, Y. Tsuchiya, Y. Fukuda, B. Zhang, T. Kuwano, A. Ike, D. Yanagi, K. Kubota, K. Mori, A. Iwata, H. Nishikawa, A. Kawamura, N. Miller, K. Matsuo, K. Shirai, K. Saku</i> . . . . .	187
Impact of CYP2C19 polymorphism on residual platelet reactivity in patients with coronary heart disease during antiplatelet therapy. <i>K. Yamamoto, S. Hokimoto, T. Chitose, K. Morita, T. Ono, K. Kaikita, K. Tsujita, T. Abe, M. Deguchi, H. Miyagawa, J. Saruwatari, H. Sumida, S. Sugiyama, K. Nakagawa, H. Ogawa</i> . . . . .	194

Elevated peripheral blood mononuclear cell count is an independent predictor of left ventricular remodeling in patients with acute myocardial infarction. <i>S. Aoki, A. Nakagomi, K. Asai, H. Takano, M. Yasutake, Y. Seino, K. Mizuno</i> . . . . .	202
Effects of mineralocorticoid receptor antagonist spironolactone on atrial conduction and remodeling in patients with heart failure. <i>M. Kimura, H. Ogawa, T. Wakeyama, A. Takaki, T. Iwami, Y. Hadano, M. Mochizuki, A. Hiratsuka, A. Shimizu, M. Matsuzaki</i> . . . . .	208
The utility of echocardiographic evaluation of donor hearts upon the organ procurement for heart transplantation. <i>S. Hashimoto, T.S. Kato, K. Komamura, A. Hanatani, K. Niwaya, T. Funatsu, J. Kobayashi, Y. Sumita, N. Tanaka, K. Hashimura, M. Asakura, H. Kanzaki, M. Kitakaze</i> . . . . .	215
Low atrial septal pacing with dual-chamber pacemakers reduces atrial fibrillation in sick sinus syndrome. <i>H. Minamiguchi, S. Nanto, T. Onishi, T. Watanabe, M. Uematsu, I. Komuro</i> . . . . .	223

## VOLUME 57, ISSUE 3 . . . . . May 2011

### Reviews

Effect of coronary endothelial function on outcomes in patients undergoing percutaneous coronary intervention. <i>S. Hamasaki, C. Tei</i> . . . . .	231
Is mechanical dyssynchrony still a major determinant for responses after cardiac resynchronization therapy? <i>Q. Zhang, C.M. Yu</i> . . . . .	239
Circadian clock and cardiovascular disease. <i>N. Takeda, K. Maemura</i> . . . . .	249
Cardiovascular disease in recent onset diabetes mellitus. <i>S. Yamagishi</i> . . . . .	257

### Original Articles

Learning curve for ablation of atrial fibrillation in medium-volume centers. <i>A. Sairaku, Y. Nakano, N. Oda, Y. Makita, K. Kajihara, T. Tokuyama, Y. Kihara</i> . . . . .	263
QT/RR relation during atrial fibrillation based on a single beat analysis in 24-h Holter ECG: The role of the second and further preceding RR intervals in QT modification. <i>A. Fujiki, R. Yoshioka, M. Sakabe, S. Kusuzaki</i> . . . . .	269
Impact of concomitant use of proton-pump inhibitors and thienopyridine derivatives on the antiplatelet effects. <i>K. Tsukahara, K. Kimura, S. Morita, T. Ebina, M. Kosuge, K. Hibi, N. Iwahashi, M. Endo, N. Maejima, T. Sugano, S. Umemura</i> . . . . .	275
The potential benefits and risks of the use of dual antiplatelet therapy beyond 6 months following sirolimus-eluting stent implantation for low-risk patients. <i>T. Nakagawa, Y. Hiasa, S. Hosokawa, T. Minami, Y. Yano, K. Yoneda, M. Mizobe, N. Murakami, Y. Tobetto, H. Chen, S. Miyazaki, K. Mahara, R. Ogura, H. Miyajima, K. Yuba, T. Takahashi, K. Kishi, R. Ohtani</i> . . . . .	283
Clinical outcomes of acute myocardial infarction with occluded left circumflex artery. <i>S.S. Kim, H.S. Choi, M.H. Jeong, J.G. Cho, Y.K. Ahn, J.H. Kim, S.C. Chae, Y.J. Kim, S.H. Hur, I.W. Seong, T.J. Hong, D.H. Choi, M.C. Cho, C.J. Kim, K.B. Seung, W.S. Chung, Y.S. Jang, S.W. Rha, J.H. Bae, S.J. Park, other Korea Acute Myocardial Infarction Registry Investigators</i> . . . . .	290
Relationship between tissue characterization with 40 MHz intravascular ultrasound imaging and 64-slice computed tomography. <i>M. Utsunomiya, H. Hara, M. Moroi, K. Sugi, M. Nakamura</i> . . . . .	297
Clinical significance of the measurements of plasma N-terminal pro-B-type natriuretic peptide levels in patients with coronary artery disease who have undergone elective drug-eluting stent implantation. <i>Y. Masaki, K. Shimada, T. Kojima, K. Miyauchi, K. Inoue, T. Kiyanagi, M. Hiki, K. Fukao, K. Hirose, H. Ohsaka, A. Kume, T. Miyazaki, H. Ohmura, A. Ohsaka, H. Daida</i> . . . . .	303
Prediction of coronary artery stenosis using strain imaging diastolic index at rest in patients with preserved ejection fraction. <i>K. Kimura, K. Takenaka, X. Pan, A. Ebihara, K. Uno, N. Fukuda, T. Kohro, H. Morita, Y. Yatomi, R. Nagai</i> . . . . .	311

Serial cardiac influence of volume overload induced by interventional therapy for central venous stenosis or occlusion in chronic hemodialysis patients. <i>Y. Horita, M. Namura, M. Ikeda, T. Tsuchiya, H. Terai, R. Fukuoka, N. Tama, T. Takagi, Y. Kumatani, Y. Ishimori</i> . . . . .	316
Serum syndecan-4 is a novel biomarker for patients with chronic heart failure. <i>R. Takahashi, K. Negishi, A. Watanabe, M. Arai, F. Naganuma, Y. Ohyama, M. Kurabayashi</i> . . . . .	325
Effect of taurine supplementation on exercise capacity of patients with heart failure. <i>M.R. Beyranvand, M. Kadkhodai Khalafi, V.D. Roshan, S. Choobineh, S.A. Parsa, M.A. Piranfar</i> . . . . .	333
Altered microRNA expression associated with reduced catecholamine sensitivity in patients with chronic heart failure. <i>H. Funahashi, H. Izawa, A. Hirashiki, X.W. Cheng, Y. Inden, M. Nomura, T. Murohara</i> . . . . .	338
Identification of novel mutations in the $\alpha$ -galactosidase A gene in patients with Fabry disease: Pitfalls of mutation analyses in patients with low $\alpha$ -galactosidase A activity. <i>M. Yoshimitsu, K. Higuchi, M. Miyata, S. Devine, A. Mattman, S. Sirrs, J.A. Medin, C. Tei, T. Takenaka</i> . . . . .	345
Aortic stiffness is strikingly increased with age $\geq 50$ years in clinically normal individuals and preclinical patients with cardiovascular risk factors: Assessment by the new technique of 2D strain echocardiography. <i>Y. Oishi, H. Miyoshi, Y. Mizuguchi, A. Iuchi, N. Nagase, T. Oki</i> . . . . .	354

## Letter to the Editor

Mortality benefit and reduced need for repeat revascularization with statin therapy among patients undergoing percutaneous coronary intervention. <i>Z.J. Zhang</i> . . . . .	360
Author's Reply. <i>M. Nakamura</i> . . . . .	361

## Corrigendum

Corrigendum to "Investigation of optimal anticoagulation strategy for stroke prevention in Japanese patients with atrial fibrillation—The J-RHYTHM Registry study design" [J. Cardiol. 57 (2011) 95-99]. <i>H. Atarashi, H. Inoue, K. Okumura, T. Yamashita, H. Origasa, for the J-RHYTHM Registry Investigators</i> . . . . .	362
--	-----

## VOLUME 58, ISSUE 1 . . . . . July 2011

### Reviews

Frontiers of anticoagulation therapy for atrial fibrillation. <i>T. Yamashita</i> . . . . .	1
Current status and prospects of antiplatelet therapy in percutaneous coronary intervention in Japan: Focus on adenosine diphosphate receptor inhibitors. <i>H. Ogawa, S. Hokimoto, K. Kaikita, K. Yamamoto, T. Chitose, T. Ono, K. Tsujita</i> . . . . .	6

### Original Articles

Risk for atrial fibrillation in patients with hypertrophic cardiomyopathy: Association with insulin resistance. <i>Y. Shigematsu, M. Hamada, T. Nagai, K. Nishimura, K. Inoue, J. Suzuki, A. Ogimoto, J. Higaki</i> . . . .	18
Restenosis after implantation of sirolimus-eluting stent begins suddenly, shows short term progression, and stops suddenly. <i>K. Tanaga, K. Miura, T. Kageyama, Y. Nakamura, T. Inoue, K. Jo, T. Ishikawa, A. Miyazaki</i> . . . . .	26
Long-term clinical outcomes after deferral of percutaneous coronary intervention of intermediate coronary stenoses based on coronary pressure-derived fractional flow reserve. <i>T. Misaka, H. Kunii, H. Mizukami, N. Sakamoto, K. Nakazato, Y. Takeishi</i> . . . . .	32
Pentraxin 3 as a biomarker for acute coronary syndrome: Comparison with biomarkers for cardiac damage. <i>N. Kume, H. Mitsuoka, K. Hayashida, M. Tanaka</i> . . . . .	38

Patients with more coronary yellow plaques have higher risk of stenosis progression within 7 months. Y. Omori, Y. Ueda, K. Okada, M. Nishio, A. Hirata, M. Asai, T. Nemoto, K. Matsuo, K. Kashiwase, K. Kodama . . . . .	46
Systolic blood pressure at admission, clinical manifestations, and in-hospital outcomes in patients with acute myocardial infarction. J. Shiraishi, Y. Kohno, T. Sawada, D. Ito, M. Kimura, M. Ariyoshi, A. Matsui, M. Arihara, H. Irie, M. Hyogo, T. Shima, T. Nakamura, S. Matoba, H. Yamada, A. Matsumuro, T. Shirayama, M. Kitamura, K. Furukawa, H. Matsubara . . . . .	54
Non-fatal cardiovascular outcome in patients with posttraumatic stress symptoms caused by myocardial infarction. R. von Känel, R. Hari, J.P. Schmid, L. Wiedemar, E. Guler, J. Barth, H. Saner, U. Schnyder, S. Begré . . . . .	61
Improvement effect on endothelial function in patients with congestive heart failure treated with cardiac resynchronization therapy. K. Enomoto, H. Yamabe, K. Toyama, Y. Matsuzawa, M. Yamamuro, T. Uemura, K. Morihisa, S. Iwashita, K. Kaikita, S. Sugiyama, H. Ogawa . . . . .	69
Change in cardio-ankle vascular index by long-term continuous positive airway pressure therapy for obstructive sleep apnea. M. Kato, T. Kumagai, R. Naito, K. Maeno, S. Kasagi, F. Kawana, S. Ishiwata, K. Narui, T. Kasai . . . . .	74

## VOLUME 58, ISSUE 2 . . . . . September 2011

### Editorial

The <i>Journal of Cardiology</i> receives its first impact factor. A. Hirayama, C. Tei . . . . .	83
--	----

### Reviews

Stent thrombosis of drug eluting stent: Pathological perspective. G. Nakazawa . . . . .	84
Stent thrombosis and drug-eluting stents. T. Takayama, T. Hiro, A. Hirayama . . . . .	92
Frontier of catheter ablation for atrial fibrillation. Y. Iesaka . . . . .	99

### Original Articles

Role of oral amiodarone in patients with atrial fibrillation and congestive heart failure. M. Kawabata, K. Hirao, H. Hachiya, K. Higuchi, Y. Tanaka, A. Yagishita, O. Inaba, M. Isobe . . . . .	108
Recent mortality of Japanese patients with atrial fibrillation in an urban city of Tokyo. S. Suzuki, T. Yamashita, T. Otsuka, K. Sagara, T. Uejima, Y. Oikawa, J. Yajima, A. Koike, K. Nagashima, H. Kirigaya, K. Ogasawara, H. Sawada, T. Aizawa . . . . .	116
Atrial fibrillation and flutter outcomes and risk determination (AFFORD): Design and rationale. T.W. Barrett, A.B. Storrow, C.A. Jenkins, F.E. Harrell Jr, K.F. Miller, K.M. Moser, S. Russ, D.M. Roden, D. Darbar . . . . .	124
Angiotensin II type 1 receptor blocker attenuates diabetes-induced atrial structural remodeling. T. Kato, T. Yamashita, A. Sekiguchi, T. Tsuneda, K. Sagara, M. Takamura, S. Kaneko, T. Aizawa, L.T. Fu . . . . .	131
Clinical study of the electrophysiological effects of ischemic post-conditioning in patients with acute myocardial infarctions. K. Okishige, S. Kanda, T. Shimura, M. Kurabayashi, D. Ueshima, N. Miwa, K. Sugiyama, H. Aoyagi, K. Yoshimura, H. Yanagi, K. Azegami . . . . .	137
Relationship between symptom-onset-to-balloon time and long-term mortality in patients with acute myocardial infarction treated with drug-eluting stents. Y.W. Cho, J.S. Jang, H.Y. Jin, J.S. Seo, T.H. Yang, D.K. Kim, D.I. Kim, S.H. Lee, Y.K. Cho, D.S. Kim . . . . .	143
Elevated concentrations of pentraxin 3 are associated with coronary plaque vulnerability. T. Soeki, T. Niki, K. Kusunose, S. Bando, Y. Hirata, N. Tomita, K. Yamaguchi, K. Koshiba, S. Yagi, Y. Taketani, T. Iwase, H. Yamada, T. Wakatsuki, M. Akaike, M. Sata . . . . .	151



Long-term outcomes of women with coronary artery disease following complete coronary revascularization. <i>H. Sato, T. Kasai, K. Miyauchi, N. Kubota, K. Kajimoto, T. Miyazaki, A. Nishino, K. Yaginuma, H. Tamura, T. Kojima, K. Yokoyama, T. Kurata, A. Amano, H. Daida</i> . . . . .	158
Rationale and design of the Japanese Heart Failure Outpatients Disease Management and Cardiac Evaluation (J-HOMECARE). <i>M. Tsuchihashi-Makaya, H. Matsuo, S. Kakinoki, S. Takechi, H. Tsutsui, for the J-HOMECARE Investigators</i> . . . . .	165
Impact of diabetes on muscle mass, muscle strength, and exercise tolerance in patients after coronary artery bypass grafting. <i>M. Nishitani, K. Shimada, S. Sunayama, Y. Masaki, A. Kume, K. Fukao, E. Sai, H. Yamashita, H. Ohmura, T. Onishi, M. Shioya, H. Sato, A. Shimada, T. Yamamoto, A. Amano, H. Daida</i> . . . . .	173
Dynamic changes in plasma total and high molecular weight adiponectin levels in acute heart failure. <i>T. Ohara, K. Hashimura, M. Asakura, A. Ogai, M. Amaki, T. Hasegawa, H. Kanzaki, M. Sonoda, H. Nishizawa, T. Funahashi, M. Kitakaze</i> . . . . .	181
Dynamic component of sports is an important determinant factor of heart rate recovery. <i>J. Nagashima, N. Matsumoto, A. Takagi, H. Musha, K. Chikaraishi, M. Sagehashi, R. Nakagawa, N. Ishige, R. Fujimaki, A. Akaike, R. Seo, H. Aoki, M. Murayama</i> . . . . .	191
<b>Letter to the Editor</b>	
White blood cell and peripheral blood mononuclear cell counts for the prediction of left ventricular remodeling after myocardial infarction. <i>A. Bauters, M. Fertin, N. Lamblin, F. Pinet, C. Bauters</i> . . .	197
Author's reply. <i>S. Aoki</i> . . . . .	198

## **VOLUME 58, ISSUE 3 . . . . . November 2011**

### **Review**

Gastrointestinal malignancies and cardiovascular diseases—Non-negligible comorbidity in an era of multi-antithrombotic drug use. <i>N. Ishizaka, A. Sakamoto, M. Fujishiro, R. Nagai and K. Koike</i> . . . .	199
---	-----

### **Original Articles**

Recurrence of angina pectoris after percutaneous coronary intervention is reduced by statins in Japanese patients. <i>R. Tsunoda, T. Sakamoto, S. Kojima, Y. Ogata, A. Kitagawa and H. Ogawa</i> . . . . .	208
<i>Chlamydia pneumoniae</i> heat shock protein 60 is associated with apoptotic signaling pathway in human atheromatous plaques of coronary artery disease patients. <i>H.C. Jha, P. Srivastava, H. Vardhan, L.C. Singh, A.R. Bhengraj, J. Prasad and A. Mittal</i> . . . . .	216
High-sensitivity cardiac troponin T in essential hypertension. <i>Y. Sato, E. Yamamoto, T. Sawa, K. Toda, T. Hara, T. Iwasaki, H. Fujiwara and Y. Takatsu</i> . . . . .	226
Physiological basis and clinical significance of left ventricular suction studied using echo-dynamography. <i>M. Tanaka, T. Sakamoto, S. Sugawara, H. Nakajima, T. Kameyama, H. Tabuchi, Y. Katahira, S. Ohtsuki and H. Kanai</i> . . . . .	232
Predictive importance of left ventricular myocardial stiffness for the prognosis of patients with congestive heart failure. <i>S. Watanabe, J. Shite, H. Takaoka, T. Shinke, Y. Tanino, H. Otake, D. Matsumoto, D. Ogasawara, T. Sawada, K. Hirata and M. Yokoyama</i> . . . . .	245
Scoring of late gadolinium enhancement in cardiac magnetic resonance imaging can predict cardiac events in patients with hypertrophic cardiomyopathy. <i>A. Nojiri, K. Hongo, M. Kawai, K. Komukai, T. Sakuma, I. Taniguchi and M. Yoshimura</i> . . . . .	253
Plasma metalloproteinase levels and left ventricular remodeling in hypertrophic cardiomyopathy in patients with an identical mutation. <i>H. Kitaoka, T. Kubo, M. Okawa, N. Takenaka, Y. Baba, N. Yamasaki, Y. Matsumura, T. Furuno and Y.L. Doi</i> . . . . .	261

Left atrial thrombus and prognosis after anticoagulation therapy in patients with atrial fibrillation. <i>S. Fukuda, H. Watanabe, K. Shimada, M. Aikawa, Y. Kono, S. Jissho, H. Taguchi, J. Umemura, M. Yoshiyama, T. Shiota, T. Sumiyoshi and J. Yoshikawa</i> . . . . .	266
Relationship between clinical outcomes and unintentional pulmonary vein isolation during substrate ablation of atrial fibrillation guided solely by complex fractionated atrial electrogram mapping. <i>Y. Iriki, S. Ishida, N. Oketani, H. Ichiki, H. Okui, Y. Ninomiya, R. Maenosono, T. Matsushita, M. Miyata, S. Hamasaki and C. Tei</i> . . . . .	278
Factors leading to failure to diagnose acute aortic dissection in the emergency room. <i>M. Kurabayashi, N. Miwa, D. Ueshima, K. Sugiyama, K. Yoshimura, T. Shimura, H. Aoyagi, K. Azegami, K. Okishige and M. Isobe</i> . . . . .	287
Impact of 64 multi-detector computed tomography for the evaluation of aortic paraprosthesis regurgitation. <i>M. Hara, M. Nishino, M. Taniike, N. Makino, H. Kato, Y. Egami, R. Shutta, J. Tanouchi, T. Hunatsu, K. Taniguchi and Y. Yamada</i> . . . . .	294